

Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

QSD11
1A422

Ind / 81a

✓

8
245

SODIUM FLUORALUMINATE AS A PESTICIDE FOR

CONTROLLING WESTERN SPRUCE BUDWORM

(CHORISTONEURA OCCIDENTALIS FREE.)



Received by: JYB
1 Index Branch

FOREST INSECT AND DISEASE MANAGEMENT
STATE AND PRIVATE FORESTRY
ROCKY MOUNTAIN REGION
USDA, FOREST SERVICE
11177 W. 8TH AVENUE
LAKEWOOD, COLORADO 80225

11033
SCPA

This publication reports studies involving pesticides. It does not contain recommendations for their use, nor does it imply that the uses discussed here have been registered. All uses of pesticides must be registered by appropriate State and/or Federal agencies before they can be recommended.

CAUTION: Pesticides can be injurious to humans, domestic animals, desirable plants, and fish or other wildlife--if they are not handled or applied properly. Use all pesticides selectively and carefully. Follow recommended practices of the disposal of surplus pesticides and pesticide containers

The use of trade, firm, or corporation names in this publication is for the information and convenience of the reader. Such use does not constitute an official endorsement or approval by the U. S. Department of Agriculture of any product or service to the exclusion of others which may be suitable.

245

SODIUM FLUORALUMINATE AS A PESTICIDE FOR
CONTROLLING WESTERN SPRUCE BUDWORM
(*Choristoneura occidentalis* Free.)

by

Robert D. L. Averill, Entomologist

Technical Report R2-8
December 1976

Forest Insect and Disease Management
State and Private Forestry
Rocky Mountain Region
USDA - Forest Service
11177 W. 8th Avenue
Lakewood, Colorado 80225

INTRODUCTION

Sodium fluoraluminate (Kryocide^(R)) is an inorganic stomach and contact poison insecticide which occurs naturally. Because of its low toxicity (LD_{50} = 10,000 mg/kg) and its ability to control a variety of Lepidopterous larvae, Kryocide was evaluated as a possible pesticide to use against western spruce budworm (*Choristoneura occidentalis* Free.) infesting high valued ornamental Douglas-fir, *Pseudotsuga menziesii* (Mirb.) Franco, trees.

METHODS

A roadside plot (ca 150m X 20m) was established in mid June 1976 by the author in a moderately defoliated stand of Douglas-fir, on the north side of Sugarloaf Mountain in Colorado (T1N, R72W, Sec. 26, NW $\frac{1}{4}$).

Ten Douglas-fir trees were selected for testing the effect of Kryocide on the budworm population. Two branches (40-45 cm in length) were removed from opposite sides of the midcrown at each sampling period. The branches removed from each tree were placed in a paper bag, sealed and labelled with the tree number and sample date. The branches were analyzed at the Lakewood Forest Insect and Disease Management laboratory within 48 hours of sampling. Data collected were budworm density and cm^2 of foliage. Population estimates are expressed in number of budworm larvae per 1000 cm^2 branch area.

Pre-spray sampling was done immediately prior to pesticide application on the plot. Post spray samples were collected four, seven, and 14 days after spraying. The budworm population was in the late fourth and early fifth stadia when the pesticide was applied.

Kryocide was applied at the rate of 2 lb active ingredient/100 gal water to the point of runoff. Application was with a Bean (Model 1010A) hydraulic sprayer equipped with a FMC 29 adjustable nozzle. Pressure was set at 200 psi.

150m = 164.04 yd
20m = 21.87 yd

40-45 cm = 15.75 - 17.72 in
1000 cm^2 = 155 sq.²in.

RESULTS

The pesticide application resulted in 95 percent larval mortality (Table 1) by post spray day 14. A high standard error is associated with the post spray budworm density estimates, partially because no population was present on six trees after post spray day seven.

Population reduction was calculated using the ratio of two ratios method developed by Simmons and Chen.

Table 1 Mean density estimates (per 1000 cm² foliage) of western spruce budworm with sodium fluoraluminate.

	Pre Spray	Post Spray Day 4	Post Spray Day 7	Post Spray Day 14
Treatment	48.2 \pm 21.5	8.3 \pm 7.2	0.47 \pm 0.7	0.6 \pm 1.1
Check	35.9 \pm 12.9	22.1 \pm 11.8	15.5 \pm 6.7	10.3 \pm 3.8
% Reduction		72	98	95

\pm values are standard error

The check population data are from another study conducted on the same mountain as the study reported herein.

DISCUSSION

Gusts of wind hampered the spray operation. Trees which lacked the appearance of an even coat of spray deposit were resprayed later the same morning when the winds were calm. The initial uneven application of the Kryocide would have gone undetected if the product was colorless. Its off-white color allows the applicator to insure adequate coverage. The 95 percent mortality level that occurred appears to be satisfactory for protecting high valued trees. The three percent difference in the post spray sampling is probably due to sampling variation. Populations of less than one larvae per 1000 cm² of foliage are difficult to locate unless a deliberate search effort is conducted.

Some settling of the pesticide was noted after the spray pump engine was turned off. Whether this is typical of sodium fluoraluminate is unknown to the author. However, it appears advisable to use spray equipment which will provide thorough agitation of the pesticide as was the case in this test.

RECOMMENDATION

It is recommended that registration be sought with the Environmental Protection Agency to label sodium fluoraluminate (Kryocide) for use against the western spruce budworm at the rate of 2 lb a.i./100 gal water applied to individual trees to the point of runoff with ground application equipment.

REFERENCES

- Simmons, G. A. and Chen, D. W. 1975. Approaches to Assessing Insecticide Efficacy in Spruce Budworm (Lepidoptera: Tortricidae) Control Programs. Can. Ent. 107:1205-1209.
- Thomson T. W. 1975. Agricultural Chemicals, Book 1 Insecticides. Thomson Publications Indianapolis Indiana p. 61-62.

